

QCD Radiation Effects on the $H \rightarrow WW \rightarrow \ell\nu\ell\nu$ Signal at Hadron Colliders

Fabian Stöckli*

Institute for Particle Physics, ETH Zurich, Switzerland

The discovery of a Standard Model Higgs boson is possible when experimental cuts are applied which increase the ratio of signal and background cross-sections. We present a study of the $pp \rightarrow H \rightarrow WW$ signal cross-section at hadron colliders which requires a selection of Higgs bosons with small transverse momentum. We compare predictions for the efficiency of the experimental cuts from a NNLO QCD calculation and the event generator MC@NLO and are able to predict a reliable number for the cross-section after the application of such experimental cuts.

1. INTRODUCTION

One of the core tasks of modern hadron collider experiments is the understanding of the electro-weak symmetry breaking mechanism. In the standard model this can be explained through a simple Higgs mechanism, where the additional complex scalar field acquires a non-vanishing vacuum expectation value, and leaves the theory with an additional scalar boson, the Higgs boson, which should be experimentally detectable.

Many different channels are and will be exploited in order to either restrict the allowed Higgs bosons mass range or, preferably, discover the Higgs boson. In the range where the Higgs boson is of a mass of about twice the W boson mass ($m_H \simeq 2 \times m_W \simeq 160 \text{ GeV}$) the branching ratio of the SM Higgs boson in a pair of W bosons is almost one, thus almost all produced Higgs bosons decay via this channel. It has been shown that this channel, with both W bosons decaying leptonically, is the most promising discovery channel for a SM Higgs boson in this mass range [1].

In contrast to other decay modes (e.g. $H \rightarrow ZZ \rightarrow 4\ell$), the Higgs invariant mass cannot be reconstructed in the W boson decay mode, due to the non-detectable neutrinos in the final state. In addition the signal events are overlaid by background processes (especially continuum W-pair and $t\bar{t}$ production) with much larger cross-sections. In order to increase the signal over background ratio rather severe experimental cuts have to be applied. A discovery (exclusion) of a certain Higgs mass hypothesis thus relies heavily on a precise knowledge of the expected number of Higgs events and the corresponding final state observable distributions in this hypothesis after the application of such restricting phase-space cuts.

In this note we discuss the effects of higher-order QCD corrections on the $H \rightarrow WW \rightarrow \ell\nu\ell\nu$ cross-section at the Large Hadron Collider (LHC) after the application of typical experimental cuts up to next-to-next-to-leading order (NNLO). Additionally we compare the results from perturbative calculations at fixed order with the parton-shower Monte Carlo program MC@NLO [2]).

2. SELECTION VARIABLES

Typical experimental event selections to enhance the signal to background ratio in the $H \rightarrow WW$ channel include cuts on lepton transverse momenta, missing transverse energy and the angle between the two charged

*Electronic address: fabstoec@phys.ethz.ch




Figure 1: Cumulative cross-section (left) and K -factor (right) as a function of the veto value for a Higgs mass hypothesis of $m_H = 165$ GeV for different fixed orders in perturbation theory. The bands correspond to a simultaneous variation of the renormalization μ_R and factorization μ_F scales in the range $[m_H/2, 2 m_H]$. The solid black line denotes the nominal cut of $p_T^{\text{veto}} = 25$ GeV in the *signal cuts* of [3]. The dashed lines denote the inclusive K -factors, i.e. the ratio of the cross-sections when no experimental cut is applied.

final state leptons in the plane transverse to the beam-pipe. In addition events with hard hadronic activity in the final state are rejected (jet-veto) in order to reduce the otherwise overwhelming $t\bar{t}$ contamination. The set of cuts used in the presented work (referred to as the *signal cuts*) are described in detail in [3].

In this section we discuss in some detail the higher order QCD effects up to NNLO on two of these selection variables, i.e. on the jet-veto (p_T^{veto}) and on the angular cut ($\phi_{\ell\ell}$). Effects on other selection variables are discussed in detail in [3]. To study the effect of the higher order corrections we define the cumulative cross-section σ_{cum} as

$$\sigma_{\text{cum}}(X^{\text{cut}}) = \int_0^{X^{\text{cut}}} \frac{\partial \sigma}{\partial X} dX, \quad (1)$$

with X a specific selection variable. We can then quantify the higher order effects at (N)NLO by computing the cumulative K -factor, defined as

$$K_{\text{cum}}^{(\text{N})\text{NLO}}(X^{\text{cut}}) = \frac{\sigma_{\text{cum}}^{(\text{N})\text{NLO}}(X^{\text{cut}})}{\sigma_{\text{cum}}^{\text{LO}}(X^{\text{cut}})}. \quad (2)$$

2.1. Jet Veto

The first selection variable of interest is the jet veto. In its course events are rejected if there is any hadronic jet with a transverse momentum p_T^{jet} larger than the jet veto value p_T^{veto} in the central detector region $|\eta_{\text{jet}}| < 2.5$. Jets are reconstructed using a k_T algorithm with an R -parameter of 0.4. In Fig. 1 the resulting cumulative cross-section (left) and cumulative K -factor (right) are shown as a function of the veto value p_T^{veto} .

While the jet-veto has no effect on the LO cross-section, for higher orders the cross-section decreases with decreasing cut value p_T^{veto} . For low cut values ($p_T^{\text{veto}} < 40$ GeV) the NLO and the NNLO band overlap and the NNLO scale variation uncertainty becomes small, indicating a convergence of the perturbative series.




Figure 2: Cumulative cross-section (left) and K -factor (right) as a function of the angular cut value $\phi_{\ell\ell}^{\text{cut}}$ for a Higgs mass hypothesis of $m_H = 165$ GeV and different fixed orders in perturbation theory. The solid black line denotes the nominal cut of $\phi_{\ell\ell}^{\text{cut}} = 45^\circ$ in the *signal cuts* of [3].

2.2. Angular Cut

The second selection variable we study is the angle between the charged leptons in the plane transverse to the beam-pipe. This variable is sensitive to the spin-structure of the initial state, and thus can serve to suppress the continuum WW background (which at lowest order comes from quark-like initial states) over the signal process (which is dominated by gluon-like initial states). While in the signal process this angle is rather small, for the background processes the charged leptons tend to be back-to-back in the transverse plane.

In Fig. 2 we present the cumulative cross-section (left) and K -factor (right) as a function of the cut on this variable for a Higgs mass hypothesis of $m_H = 165$ GeV. In contrast to the jet-veto cut the K -factor increases when the cut value is decreased. It can be concluded that the application of an inclusive K -factor would fail to describe the cross-section after such cuts reliably.

3. RESUMMATION EFFECTS

After having studied the fixed-order QCD corrections on some of the selection variables we are interested in comparing the results with cross-sections computed using Monte-Carlo event generators, i.e. MC@NLO, in which leading logarithmic terms are re-summed up to all orders in perturbation theory by the application of a parton shower algorithm. In order to being able to compare the MC@NLO spectra with the fixed NNLO predictions we re-weight the MC@NLO numbers (which are nominally at NLO) to the inclusive NNLO prediction. These results are labeled as $R(\text{MC@NLO})$.

We show here the comparison of the cumulative cross-sections in two of the selection variables, for the jet-veto and a cut on the missing transverse energy (E_T^{miss}). The corresponding distributions are shown in Fig. 3. While the small discrepancy between the fixed and the re-summed result in the jet-veto plot (left) can be explained by the presence of more partons in the final state in the MC@NLO case, the E_T^{miss} -distributions agree perfectly. A more detailed discussion can be found in Ref. [4].




Figure 3: Comparison between fixed-order perturbation theory and inclusively re-weighted MC@NLO of the cumulative cross-sections in the jet veto value p_T^{veto} (left) and the missing transverse momentum E_T^{miss} (right) for a Higgs mass hypothesis of $m_H = 165$ GeV. The bands correspond to a variation of the factorization and renormalization scales in the range $[m_H/2, 2 m_H]$.

4. ACCEPTED CROSS SECTION

We finally compute the cross-sections after the application of all *signal cuts* of Ref. [3] in the different orders in perturbation theory and compare them to the results from MC@NLO. The numbers are shown in the following table.

σ_{acc} [fb]	$\mu = m_H/2$	$\mu = 2 m_H$
LO	21.00 ± 0.02	14.53 ± 0.01
NLO	22.40 ± 0.06	19.52 ± 0.05
MC@NLO	18.42 ± 0.08	14.39 ± 0.06
NNLO	18.45 ± 0.54	19.01 ± 0.27
$R^{\text{NNLO}}(\text{MC@NLO})$	20.43 ± 0.09	18.24 ± 0.07

While at NLO the comparison with MC@NLO fails, the comparison between NNLO and inclusively re-weighted MC@NLO shows a very good agreement. From this we can conclude that the NNLO cross-sections in the table are a very precise prediction for the expected Higgs cross-section after the application of the *signal cuts*.

References

- [1] M. Dittmar and H. K. Dreiner, Phys. Rev. D **55**, 167 (1997) [arXiv:hep-ph/9608317].
- [2] S. Frixione and B. R. Webber, JHEP **0206**, 029 (2002) [arXiv:hep-ph/0204244].
- [3] C. Anastasiou, G. Dissertori and F. Stöckli, JHEP **0709** (2007) 018 [arXiv:0707.2373 [hep-ph]].
- [4] C. Anastasiou, G. Dissertori, F. Stöckli and B. R. Webber, JHEP **0803** (2008) 017 [arXiv:0801.2682 [hep-ph]].